

JOSE I. ARAUZ, DMD, PA

PRACTICE LIMITED TO PERIODONTICS AND DENTAL IMPLANTS DIPLOMATE OF THE AMERICAN BOARD OF PERIODONTOLOGY 200 Doctors Drive, Suite N Jacksonville, NC 28546

(910) 577-1315 main

fax (910) 577-1078

PRESCRIPTIONS FOR SURGERY

WE ADVISE TAKING ALL MEDICATIONS WITH FOOD. THERE ARE NO DIETARY RESTRICTIONS PRIOR TO SURGERY.
THE FOLLOWING MEDICATIONS ARE TO REDUCE INFLAMMATION AND AID HEALING:
MEDROL 5 MG - 21 TABLETS - (TAKE AS DIRECTED BELOW, TO START DAY BEFORE SURGERY)
PREDNISONE 5 MG - 21 TABLETS - (TAKE AS DIRECTED BELOW, TO START THE DAY BEFORE SURGERY) THESE SHOULD BE TAKEN AS FOLLOWED: 1ST DAY TAKE 6 TABLETS: 2 AT BREAKFAST / 2 AT LUNCH / 2 AT DINNER 2ND DAY TAKE 5 TABLETS: 2 AT BREAKFAST / 2 AT LUNCH / 1 AT DINNER 3RD DAY TAKE 4 TABLETS: 2 AT BREAKFAST / 1 AT LUNCH / 1 AT DINNER 4TH DAY TAKE 3 TABLETS: 1 AT BREAKFAST / 1 AT LUNCH / 1 AT DINNER 5TH DAY TAKE 2 TABLETS: 1 AT BREAKFAST / / 1 AT DINNER
6 th DAY TAKE 1 TABLET: 1 AT BREAKFAST THE FOLLOWING ANTIBIOTICS MAY BE PRESCRIBED FOR GUM SURGERY, BONE GRAFTING, IMPLANTS, AND OTHER SURGICAL PROCEDURES. THERE IS NO NEED TO TAKE A PREMED FOR JOINT REPLACEMENT ON THE DAY OF SURGERY: AMOXICILLIN 500 MG — (TAKE 1 CAPSULE 3 TIMES DAILY UNTIL FINISHED, TO START DAY BEFORE SURGERY)
AUGMENTIN MG – (TAKE 1 TABLET 2 TIMES DAILY UNTIL FINISHED, TO START DAY BEFORE SURGERY)
AZITHROMYCIN Z-PACK 250 MG (TAKE AS DIRECTED, TO START DAY BEFORE SURGERY)
AZITHROMYCIN 250 MG –(TAKE 2 TABS ON FIRST DAY, THEN 1 TAB DAILY UNTIL FINISHED, TO START DAY BEFORE SURGERY)
CLINDAMYCIN 300 MG – (TAKE 1 CAPSULE 3 TIMES DAILY UNTIL FINISHED, TO START DAY BEFORE SURGERY)
METRONIDAZOLE 500 MG— (TAKE 1 TABLET 3 TIMES DAILY UNTIL FINISHED, TO START DAY BEFORE SURGERY)(TAKE;TIMES DAILY UNTIL FINISHED, TO START DAY BEFORE SURGERY)
ONE OF THE FOLLOWING MEDICATIONS WILL BE PRESCRIBED FOR PAIN FOLLOWING PROCEDURE:
ULTRACET – (TAKE 2 TABS EVERY 4-6 HOURS,TO START FOLLOWING SURGERY) for the first 3 days; then as needed for pain
IBUPROFEN 800 MG — (TAKE 1 TABLET EVERY 6 HOURS, TO START FOLLOWING SURGERY) for the first 3 days; then take 1 tab every 8 hours as needed for pain
VICODIN 5/300 MG – (TAKE 2 TABLETS EVERY 6 HOURS, TO START FOLLOWING SURGERY) as needed for pain
PERCOCET 5/325 MG — (TAKE 2 TABLETS EVERY 6 HOURS, TO START FOLLOWING SURGERY) as needed for painNORCO 5/325 MG — (TAKE 1 or 2 TABLETS EVERY 6 HOURS, TO START FOLLOWING SURGERY) as needed for pain(TAKE;TIMES DAILY, TO START FOLLOWING SURGERY)
AN ANTIBACTERIAL MOUTHRINSE OR GEL WILL BE PRESCRIBED TO KEEP THE SURGERY AREA CLEAN DURING THE INITIAL HEALING PHASE:
CHLORHEXIDINE ORAL RINSE 0.12% 16 OZ — (RINSE WITH 15 ML 2 TIMES DAILY, ONCE AFTER BREAKFAST AND ONCE BEFORE BEDTIME, TO START DAY BEFORE SURGERY)